


Ages For Exoplanet Host Stars


Rotation & Lithium in the Kepler Field


Luke Bouma

Carnegie Fellow · Observatories


ExSoCal · December 15, 2025









Most exoplanets







Most exoplanets




Radius valley







Most exoplanets




Radius valley




Compact multis







Most exoplanets



Radius valley



Compact multis



$\eta_{\oplus}$



Where are Kepler's missing young planets?

Where are Kepler's missing young planets?

5,000 planets. $t/\text{Gyr} \sim \text{U}[0,10]$

Age distribution:

Binney+00, Nordstrom+04

Where are Kepler's **missing young planets**?


5,000 planets. $t/\text{Gyr} \sim \text{U}[0,10] \Rightarrow$ Expect 500 sub-gigayear planets.

Age distribution:


Binney+00, Nordstrom+04

Where are Kepler's **missing young planets**?



5,000 planets. $t/\text{Gyr} \sim \text{U}[0,10] \Rightarrow$ Expect 500 sub-gigayear planets. 50 observed!






Sun-like stars spend gigayears on main seq.




**Sun-like stars spend
gigayears on main seq.**



**Sun-like stars spend
gigayears on main seq.**



Stellar ages: open clusters anchor rotational ages for field stars




Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gillen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rappelli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20

Stellar ages: open clusters

anchor rotational ages for field stars




Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gillen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rappelli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20

Stellar ages: open clusters anchor rotational ages for field stars




Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gillen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rampalli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20


Stellar ages: open clusters anchor rotational ages for field stars



Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

Stellar ages: open clusters anchor rotational ages for field stars




Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gillen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rappelli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20

Stellar ages: open clusters anchor rotational ages for field stars




Binaries are removed;
Fit from 3800 to 6200 K

Bouma+23

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gillen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rappelli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20

Stellar ages: open clusters anchor rotational ages for field stars





Binaries are removed;
Fit from 3800 to 6200 K



 `pip install gyro-interp`

α Per: Boyle & Bouma 23, Pleiades: Rebull+16, Blanco-1: Gilen+20, Psc-Eri: Curtis+19b, NGC-3532: Fritzewski+19
Group-X: Messina+22, Praesepe: Rappelli+21, NGC-6811: Curtis+19a, NGC-6819: Meibom+15, Ruprecht-147: Curtis+20

Kepler detected rotation for $\approx$ one in three of its stars




Lithium in Keck/HIRES spectra of KOIs provides independent ages




Model: **Jeffries+23**, Weaver+24
Li: Fulton+17, Johnson+17, Petigura+17, Berger+18, Bouma+24


Kepler's open clusters provide extra verification




Young Kepler stars **3-5x more rare** than their 3 Gyr counterparts



Young Kepler stars **3-5x more rare** than their 3 Gyr counterparts









Kepler

$(l, b) = (77, +13^\circ)$
 $\langle z \rangle = 230 \text{ pc}$


Kepler's age distribution is a product of both kinematic heating & star formation history



Field ages offer **2-3x expansion** for demographics at 0.1-1 Gyr



Field ages offer **2-3x expansion** for demographics at 0.1-1 Gyr



Precision limit for rotational & lithium ages set by **intrinsic scatter** in open cluster distributions.

- Rotation: **gyro-interp** (Bouma+23)
- Lithium: *eagles* (Jeffries+23, Weaver+24)
- Calibrated for: single FGK stars at < 4 Gyr near solar metallicity

Youngest Kepler stars **3-5x rarer** than stars 3 Gyr old.

- Age distribution product of both **kinematic heating** and **declining SFR**
- Unresolved binaries main source of uncertainty

Kepler's missing young planets **no longer missing**.


- 10x discrepancy now $< 2x$, due to **declining dN/dt** and **new rotational ages**
- $N=109$ Kepler planets with $t < 1$ Gyr: useful for demographics (Bouma+24)
- TESS, PLATO, Roman, & Rubin will open new stellar populations, young & old

Luke Bouma

lbouma@carnegiescience.edu

bonus

How precise are gyro ages?



1) Interpolates between clusters

0.08 - 4 Gyr, FGK (3800-6200 K)


2) Fits the scatter

Normal + trailing uniform distribution

Let uniform distribution converge with ZAMS

3) Calculates $p(t|P_{\text{rot}}, T_{\text{eff}})$ by marginalizing over scatter

.....



1) Interpolates between clusters
0.08 - 4 Gyr, FGK (3800-6200 K)


2) Fits the scatter
Normal + trailing uniform distribution
Let uniform distribution converge with ZAMS

3) Calculates $p(t|P_{\text{rot}}, T_{\text{eff}})$ by
marginalizing over scatter


► Reproduces <4 Gyr spin-down
With the evolving fast fraction!
With stalled spin-down!



► $P_{\text{rot}}, T_{\text{eff}} \nRightarrow$ unique age
Range of ages dictated by scatter in data

 `pip install gyro-interp`



How do unresolved binaries affect gyro ages?





gyro-interp (Prot) vs. eagles (Li) age scale comparison

They mostly* agree, but not many stars overlap!

Bouma+24 suppl.

How good is $P_{\text{rot}} \sim t^{1/2}$?

